

02-22-00

A

## FISH &amp; RICHARDSON P.C.

225 Franklin Street  
Boston, Massachusetts  
02110-2804

Telephone  
617 542-5070

Facsimile  
617 542-8906

Web Site  
www.fr.com

February 18, 2000

Attorney Docket No.: 04930-028001

**Box Patent Application**

Assistant Commissioner for Patents  
Washington, DC 20231

Presented for filing is a new original patent application of:

Applicant: HUGH S. KEEPING and JONATHAN S. REICHNER

Title: TREATMENT FOR BONE DISORDERS

Enclosed are the following papers, including those required to receive a filing date  
under 37 CFR 1.53(b):

	<u>Pages</u>
Specification	21
Claims	4
Abstract	1
Declaration	[To be Filed at a Later Date]
Drawing(s)	5

Enclosures:

— Postcard.

## CERTIFICATE OF MAILING BY EXPRESS MAIL

Express Mail Label No. EL228030161US

I hereby certify under 37 CFR §1.10 that this correspondence is being  
deposited with the United States Postal Service as Express Mail Post  
Office to Addressee with sufficient postage on the date indicated below  
and is addressed to the Assistant Commissioner for Patents, Washington,  
D.C. 20231.

Date of Deposit February 18, 2000Signature *Lisa G. Gray*

Lisa G. Gray  
Typed or Printed Name of Person Signing Certificate

02/18/00  
JC690 U.S. PTO

Frederick P. Fish  
1855-1930

W.K. Richardson  
1859-1951

BOSTON

DELAWARE

NEW YORK

SILICON VALLEY

SOUTHERN CALIFORNIA

TWIN CITIES

WASHINGTON, DC

jc542 U.S. PTO  
09/507239  
02/18/00

FISH & RICHARDSON P.C.

Assistant Commissioner for Patents  
February 18, 2000  
Page 2

This application is entitled to small entity status. A small entity statement will be filed at a later date.

Basic filing fee	\$345
Total claims in excess of 20 times \$9	\$162
Independent claims in excess of 3 times \$39	\$78
Fee for multiple dependent claims	\$0
Total filing fee:	\$585

A check for the filing fee is enclosed. Please apply any other required fees or any credits to deposit account 06-1050, referencing the attorney docket number shown above.

If this application is found to be incomplete, or if a telephone conference would otherwise be helpful, please call the undersigned at (617) 542-5070.

Kindly acknowledge receipt of this application by returning the enclosed postcard.

Please send all correspondence to:

INGRID A. BEATTIE, PH.D., J.D.  
Fish & Richardson P.C.  
225 Franklin Street  
Boston, MA 02110-2804

Respectfully submitted,



Ingrid A. Beattie, Ph.D., J.D.  
Reg. No. 42,306  
Enclosures  
IAB/lgg  
20029335.doc

APPLICATION  
FOR  
UNITED STATES LETTERS PATENT

TITLE: TREATMENT FOR BONE DISORDERS

APPLICANT: HUGH S. KEEPING AND JONATHAN S. REICHNER

CERTIFICATE OF MAILING BY EXPRESS MAIL

Express Mail Label No. EL228030161US

I hereby certify under 37 CFR §1.10 that this correspondence is being deposited with the United States Postal Service as Express Mail Post Office to Addressee with sufficient postage on the date indicated below and is addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231.

February 18, 2000

Date of Deposit

Signature

Typed or Printed Name of Person Signing Certificate

## TREATMENT FOR BONE DISORDERS

### BACKGROUND OF THE INVENTION

The invention relates to bone loss disorders. Catastrophic bone loss, i.e., osteolysis,  
 5 is a debilitating pathological consequence of a spectrum of disease states including  
 rheumatoid arthritis, osseous metastasis, aseptic prosthetic loosening and periodontitis.  
 Rheumatoid arthritis (RA) is a chronic inflammatory disease which often results in long term  
 disability and increased mortality.

### SUMMARY OF THE INVENTION

10 The invention provides compositions and methods to deliver an anti-inflammatory  
 composition, e.g., recombinant human interleukin-4 (rhIL-4), to build (or rebuild) bone  
 tissue. The composition is produced from living osteoprogenitor cells (OPCs) or  
 odontoprogenitor cells. The cells contain a genetically-engineered viral or non-viral plasmid  
 vector containing a regulatable, inducible, osteoblast-specific promoter to direct expression  
 15 of an anti-inflammatory polypeptide at specific sites of implantation in bone to inhibit  
 osteolysis. For example, a bone stromal cell is isolated from autologous or allogeneic  
 periodontal ligament and manipulated *ex vivo* prior to implantation into a recipient patient.  
 Stromal cells are cultured in the presence of extracellular matrix (ECM) components to  
 differentiate into odontoprogenitor cells. For example, ECM contains bone morphogenic  
 20 proteins (BMPs) such as BMP-6. Induction of differentiation to progenitor cells is carried  
 out before or after genetic manipulation of the cells.

Preferably, the nucleic acid with which the cells are transfected or transduced encodes  
 an anti-inflammatory cytokine or anti-inflammatory fragment of the cytokine. For example,  
 the cytokine is interleukin-4 (IL-4). The nucleic acid encodes a polypeptide containing the  
 25 amino acid sequence of SEQ ID NO:1; for example, the nucleic acid contains the coding  
 region of the nucleotide sequence of SEQ ID NO:2.

Table 1: Human IL-4 Amino Acid Sequence

MGLTSQLLPPLFFLLACAGNFVHGHKCDITLQEIIKTLNSLTEQKTLCTELTVTDIFAA  
SKNTTEKETFCRAATVLRQFYSHHEKDTRCLGATAQQFHRHKQLIRFLKRLDRNLW  
GLAGLNSCPVKEANQSTLENFLERLKTIMREKYSKCSS (SEQ ID NO:2, GENBANK™  
Accession No. M13982)

Table 2: Human IL-4 Nucleotide Sequence

1 gatcgttagc ttctcctgat aaactaattg cctcacattg tcaactgcaaa tcgacaccta  
61 ttaatgggtc tcacctccca actgcttccc cctctgttct tctgtctagc atgtgccggc  
121 aactttgtcc acggacacaa gtgcgatatc accttacagg agatcatcaa aactttgaac  
181 agcctcacag agcagaagac tctgtgcacc gaggtagccg taacagacat cttgtctgcc  
241 tccaagaaca caactgagaa ggaaaccttc tgcagggtcg cgactgtgct ccggcagttc  
301 tacagccacc atgagaagga cactcgtctc ctgggtgcga ctgcacagca gttccacagg  
361 cacaagcagc tgatccgatt cctgaaacgg ctgcacagga acctctgggg cctggcgggc  
421 ttgaattcct gtcctgtgaa ggaagccaac cagagtacgt tggaaaactt cttgaaagg  
481 ctaaagacga tcatgagaga gaaatattca aagtgttga gctgaatatt ttaattatg  
541 agttttgat agctttatt ttaagtatt tatatattta taactcatca taaaataaag  
601 tatatataga atct SEQ ID NO:2, GENBANK™ Accession No. M13982; coding  
sequences span nucleotides 64-525; signal peptide encoded by nucleotides 64-135).

Alternatively, the cells contain a nucleic acid encoding an IL-4 fragment, agonist or mutant. The fragment, agonist or mutant has anti-inflammatory activity. For example, the mutant contains a mutation in the region of IL-4 which is involved in binding to IL-2R gamma, e.g., Arg 21 is changed to a Glu residue. Sequences which differ from the coding sequence of SEQ ID NO:2 hybridize under stringent conditions, with all or part of the reference sequence and encode an anti-inflammatory polypeptide. Promoter or transcriptional regulatory elements which differ from a reference sequence hybridize under stringent conditions to a nucleic acid having the reference sequence and retain transcription regulatory function, e.g., cell specificity, of the reference sequence. For example, the nucleic acid may contain one or more sequence modifications in relation to a reference sequence. Such modifications may be obtained by mutation, deletion and/or addition of one or more nucleotides compared to the reference sequence. Modifications are introduced to alter the activity of the regulatory sequence, e.g., to improve promoter activity, to suppress a transcription inhibiting region, to make a promoter constitutive or regulatable or vice versa. Modifications are also made to introduce a restriction site to facilitate subsequent cloning

steps, or to eliminate the sequences which are not essential to the transcriptional activity. Preferably, a modified sequence is at least 70% (more preferably at least 80%, more preferably at least 90%, more preferably at least 95%, more preferably at least 99%) identical to a reference sequence. The modifications do not substantially alter the biological function of a polypeptide or the cell-specificity of transcription promoter function associated with the reference sequence.

Nucleotide and amino acid comparisons are carried out using the Lasergene software package (DNASTAR, Inc., Madison, WI). The MegAlign module used was the Clustal V method (Higgins et al., 1989, CABIOS 5(2):151-153). The parameters used were gap penalty 10, gap length penalty 10.

Alternatively, nucleic acids which differ from a given reference sequence hybridize at high stringency to a strand of DNA having the reference sequence, or the complement thereof. Hybridization is carried out using standard techniques, such as those described in Ausubel et al. (Current Protocols in Molecular Biology, John Wiley & Sons, 1989). "High stringency" refers to nucleic acid hybridization and wash conditions characterized by high temperature and low salt concentration, i.e., hybridization at 42 degrees C, and in 50% formamide; a first wash at 65 degrees C, 2 × SSC, and 1% SDS; followed by a second wash at 65 degrees C and 0.2% × SSC, 0.190 SDS. Lower stringency conditions suitable for detecting DNA sequences having about 50% sequence identity to a reference gene or sequence are detected by hybridization at 42 degrees C in the absence of formamide; a first wash at 42 degrees C in 6 × SSC and 1% SDS; and a second wash at 50 degrees C in 6 × SSC and 1% SDS.

A heterologous nucleic acid encoding a polypeptide (e.g., IL-4) is operably linked to an osteoblast-specific promoter such as an osteocalcin promoter sequence (e.g., a nucleic acid containing the nucleotide sequence of SEQ ID NO:3) or a bone sialoprotein promoter sequence (nucleotides 1-2472 of SEQ ID NO:4) or dentin sialoprotein promoter sequence (SEQ ID NO:6 and/or 7).

Table 3: Human Osteocalcin Regulatory Region

1 ttctcctgtc cggatgcgca gggcagggct gaccgtcgag ctgcacccac agcaggctgc  
 61 ctttggtgac tcaccgggtg aacgggggca ttgcgaggca tcccctccct gggtttggct  
 121 cctgcccacg ggccgtgacag tagaaatcac aggcgtgag acagctggag ccagctctg  
 5 181 ctgaacctt ttttaggtct ctgatcccg ctctcttt agactccct agagctcagc  
 241 cagtgtcaa cctgaggctg ggggtctctg aggaagagt agttggagt gaggggtctg  
 301 gggctgtccc ctgagagagg ggccagaggc agtgtcaaga gccgggcagt ctgattgtgg  
 361 ctacccctcc atactccca ggggcccctg gccagcagc cgcagctccc aaccacatat  
 421 cctctgggtg ttggcctacg gagctggggc gcatgacccc caaatagccc tggcagattc  
 10 481 cccctagacc cgccgcacc atggtcaggc atgcccctcc tcatcgctgg gcacagccca  
 541 gagggataa acagtgtctg aggcgtggcg ggcaggccag ctgagtctg agcagca  
 (SEQ ID NO:3; , GENBANK™ Accession No. E13404)

Table 4: Mouse bone sialoprotein promoter region and cDNA

1 tctagaaagc actgttctt taaaatcatt caccacctt ggctcctaca atcttctgt  
 15 61 cctccctcc acacagatcc ctgagcctt aggagagggc tgtataaat catcccctt  
 121 ggagttagca gtctgaagtc tctattctc catgactgt ctattccgt ccgcgggat  
 181 tcagttattc gtgggtgca gggggaccac gaacctggaa ggaaatggga ggaagaaaa  
 241 gagagcggac gaccaagtag attgaacata tcaaggtctc gtttattagg ctgaggtgcc  
 301 ttcttttaa agcatacatc acggggaata tgggaggggt cgaggagaa ttatacaaag  
 20 361 aacaaagaag tgggcatctg ctgacatggg ggccgaagtc aggcgccagg cagcgggcac  
 421 tctggatttt atctctggaa cattgatcct cctgacagc ctgggggtc aggctgggt  
 481 caggcgtaac tcatgtctt ggatggcagc ggaactcagg aagagatagg gaagagggga  
 541 ctataattca gctttacag cctcaggtgc caagaaagga ataggagga aggggggtga  
 601 taaccagctc ttagtacaag gccattggc ctgttagga gattgtgaag ggctcattt  
 25 661 ctacgggat ggtctctgac actgtctgc tgtgtctc ccatctact gcaagactgg  
 721 gctttctga tgaagttaa gcctagttag ggtgccctgt tcatagaag tcatattgca  
 781 gtactcagc agaatattag tagtgggtt tttccctc gagagctcac aacctgtcta  
 841 gtctgggtt ctagaccg tgaataatt tatttcaga agttaacatc ctcccctca  
 901 gacaccttg aagctgtgg gtgttgggt ttctgtccc tctacctgca cgtcttcca  
 30 961 taccacactg tgagcatttg aaagcgtgtg ctagagttc ttgttagct cccatgtcc  
 1021 tataaaacac ttggtttg tagagaactg agcagttca acttggctca actgagctta  
 1081 tgggggtgaa ttgaatacaa gcaataaaa ggagctatt caacttctt ttgtgggtc  
 1141 tctatttat tttaaatgc tgaataactt ttcttagct aaatcatctg aagaatctaa  
 1201 cagagtcact actctggcaa caatactgga caacaatggc atttattgat ttctgtaaag  
 35 1261 tagaagtcaa cagagaagaa tatggggata aagaatatag ggataaagaa gacaaccaac  
 1321 cagagctccc aggtctaaa ccaccaacca gggagtacac atggaggac ccatggctcc

1381 atctgtatat gtagcagagg atggcctagt ccatcatcaa tgggatgaga ggcccttgg  
1441 cccatgaagg cctgatatcc cagtgtcggg gaatttgagg gcagggagga gagagtggat  
1501 gggtaggtgg gggaacaccc tcatagaagc aggagggggg gtgggatagg gggtttggg  
1561 gtgtgggaat tgggaaagg gataaacctt gaaacgtaaa taaataaaat atccaataaa  
5 1621 aaaatcttct ggaaaagaaa agatatacaa aatacaaagg cagtttctt tgcaaaccta  
1681 ggaaatgttc agtttgccaa tgcattgcagt aagtttatt tccagtaatt attcaataac  
1741 catgaactgc tctctggcag tgctagtaat tattctctac tcataggaaa aaaattacat  
1801 aagaagacga ctgaaataa gattatacga tgtgcagtgg cctcattac acagcaaagg  
1861 gccacatagg ggataatccc aaggactgt tctatgaaag gttacatcag ctcttggtc  
10 1921 tcaacctcga acgctgtaac gttcacagtc agcattgtgc ttagcaaag cttaggtaat  
1981 ctgactggtt taataatct agtttgact tacaagcctc tgaaatatgt tttagggaga  
2041 aatataaagg aatcaatatt aaactatctc ttggcatcaa ctcatctct aattcagtac  
2101 tttagaccc atgcagtgtc gtgtgaaagc cagcttctct tctttcaac acagtgaana  
2161 cctgtatcat tgtgaaagct taaatgcta agtctttgc tattatttt attgaaatg  
15 2221 cagtatatta ttatatatat tcagaactct aactaccatc ttctctcac cttcaatta  
2281 aatcccacaa tgcaagcctc ttggcagaag gccaccttt catgtttatt caactgaggc  
2341 tgaatcttga aaatgtgtt agtttggga ttctctgtg agaaccaca gcctgacgtt  
2401 gtgctggcca cagctgtgat tggctgtga gaggcggaga agggtttata gtcagcaaga  
2461 gcaagtgaat gagtgagtga cagccgggag aacaatccgt gccactact cgactcgagc  
20 2521 caaggacctg gccgaaagga aggttaaggt aatgggcaag gacctcacag ccaggtaatg  
2581 ggcaaggacc tcacagccag gcacctcagt ctccctgtg tggcttggc ttggagttg  
2641 tagctgcagc atggatctta ctgcacagtg cacagtggct ctagtgaac tttagctgc  
(SEQ ID NO:4; GENBANK™ Accession No. AF071079; promoter region,  
nucleotides1-2472)

Table 5: Rat bone sialoprotein promoter region

1 aagcttaggg aacattcagc ctgccaacat acgcgggaag ttattttcc agtgatcctt  
61 tcaatggccg tggaactgct ttctggcagt gctagtaatt ctctctct cagagggaaa  
121 gatacatagg aagaggactt agaaataagc ctgagagtat acagcgcttg atgacctcac  
30 181 tcgcacaacg aaaggccatg tcccgatga tgccaactac ttgttcgat gagagttaa  
241 tcagcttctt ggtctgagcc tcaaatgtt tagctttcac agtcagcaca gtagcaaa  
301 ccttggcagc ccggctggct ttacaatact gattctgact tacgagcctc tgaaatgcat  
361 ttcaaaaagg aatataaagg gatcttact gaacacctct tgcatacaac tctttccta  
421 attcagtgtt ttagggtcg gccagtgtg tgttaacag aggcagttt tctttctt  
35 481 caacatagta aaaacctgta tcatttgtaa agtttaaatg ctaagtctg ttgccattta  
541 gtttatttga aatgcagtgt attattatag atattcagaa cttaactac catcttctc



601 tcagccttca attaaatccc acaatgcgac ctcttggcag caggcgcgcc ttcatgttt  
661 attcaactga ggctgagtct tgaacacgtg ttgtagtac ggattttctg gtgagaaccc  
721 acagcctgac gtcgcaccgg ccgtgaccgt gattggctgc tgagaggaga agaagggttt  
781 ataggctcag aagagcgagt gaatgggtga gaggcagccg ggagaacaat ccgtgccact  
5 841 cactcacttg ctctctccag ccaggactgc cgaaggtaag gtaatgggcc agcacctcac  
901 agccacctgc ctgaggcttc ctgtgtggct ttggcttga atttgtgtt gaagcatgga  
961 tcttactgct tgggtcacaa tggctctggt tgaacttag ctgtctgtga aatgggacct  
1021 ctgagtttag gttcttcca aagaccaggc tgggtaacgt aagcatgcag ttaactgct  
1081 tcagattggt acc  
10 (SEQ ID NO:5; GENBANK™ Accession No. LO6562)

Table 6: Mouse dentin sialoprotein gene regulatory regions (5' to exon 1)

1 gaattotttt cccattggta acgtaaaaga ccactactta attgagttag cttaggtcga  
15 61 acaaacagac ttatataac ttaacttctt tcacattat gaaaaattaa tcagtatcgg  
121 cactgagaag gcagaaacag gtgaaactcc atgagtttca ggccagcctg atctacatag  
181 gaattctagg acaagcaggg ctaggtagag ataccctatc tcaaaaaacc aaaacccaaa  
241 aacattacgt ttaagcagat ttagtttga cctaaatgt ttgtcttagt gaaggtccca  
301 aatgctctta gcaaatgttt cttgtgtag ttggagagtg ttgtgtgcta atacagctat  
20 361 caagcacttc ttgttagaca ccgaagatct tcttaactct ccacaggtc tggagagctg  
421 ttcaaatctg ctattacaac caagtttaga agaggaaggc aattcctgag gaaagtggca  
481 ttctaaata tgattggccc ttaagatgc tcaagaacc aagaacctg cagtgtaaat  
541 aatagcaaag tgtttactat ggaagtgcag cttcgaggaa actcccttc tatcactgga  
601 acctgtcaa tccctaccta catgaatat ttgttaatt ctctcagat aaagctctga  
25 661 agatgctgtt gctggatagt gatttaatat ttctgatcat atgtgttga catctttcag  
721 tagtgtgaca taaaacatg gacacatccc taagctggta cacagagact ccaattgcct  
781 agtgtggagc tcataagcta gagaaatggc tcagggatca tctgtatat ccagggtcag  
841 agagaatgat ggggtcaggc aagtactttt tctttctgg aagcacagcc tgttttcta  
901 ttctgtactc tatagtttac acatatagtg gagcaaagaa tgaaagctgt gtctgtggtg  
30 961 tgtgtgtgtg tgcactctgt acttacgcat agataccta caccatgtt caccttggga  
1021 acagctattt ttaaatttag ttgtattaa attaatagat tataaagaaa aacccaaaac  
1081 cttatgtca gtgttagat taaatcagaa aggtttcctg aagtactgt ttataaattc  
1141 ttttaaagat cccttaggca gtgtcaagac tgttgcagc ggacagccgc ttgaattata  
1201 gcgcaccaac ttaatatgt acctcaggaa tgataggggt cttaaatagc cagtcgtatt  
35 1261 tactagagaa acctagagtt ttcttagatt gccgacctaa gcaagaggag aaatgcaggg  
1321 tgacagagtc taagtggctc tttcagata tatcacactg attatctata ttaagacac  
1381 aaaacagtct tccaggagct atttaattaa gtgaaagtaa gtctagtct tttggaacca  
1441 aagggtctag tgagccaacg tacggcgag cgagggagtg gggcggtatt acagcctcat

1501 aggcacactg actctttaa ccccccacac agggatccta agcagtgatt ggttgagaaa  
 1561 attatcaaac tgaatttaa tttagcagg tacaaaattg tcacgcaaaa agcccaggac  
 1621 agtgtgc (SEQ ID NO:6; nucleotides 1-1627 of GENBANK™ Accession No.  
 AJ002141).

Table 7: Mouse dentin sialoprotein gene regulatory regions (intron between exon 1 and exon 2)

	gtaagat ggactccctc ctgccaggag ccaactgtct cctgttgaga
	1741 gaatctccag ctgcagagat gaggggtgact tgggataaag ttttaactc ttcaggctca
10	1801 cactatata taaagataat gtgtgattca ggaaggggtg ctaagccatc tgatgagacc
	1861 atctgataag acgacgaatc actggggagc agaactgatt ttgccccagt atattgttga
	1921 gactttaict cctataggaa aaacctaga tgaacaaac attctaattg tattaattaa
	1981 aaaaaaacag tacctgaagg gtttatgta tagttctcta tagctctatt ttgttattt
	2041 tcattcagga aaatactttt aagagctata aacctagtca aagggtgttt acagcctgtg
15	2101 ccttggaaatg ttgggagtgt tgggatttaa caaatgagaa tcacacactg tcttctctt
	2161 cgagacagag acatggatga tgcagtgtcc aaacaccagc tcttctgaa aaataagctg
	2221 ggtttggggg ttgattttaa tcatggctct tcatgatttc aaggctgcc tagtgtttat
	2281 gattaaagct ctatggcga aagaattgtg gtctctcca gggctcagta tctgctgat
	2341 attaacctcc gatgttact gactggacct aataaataaa tctcattta aacttagtat
20	2401 ctgactcag agtcaactta ggaatctgga gcgtaattt ctggcatgtg atgtgaagt
	2461 tctaaaagta gacgtcaaa cagtttatg tagaaaacac acagatctgt caagctgatt
	2521 tttagctcc aaatttcag ataataaggt tagggaaaac aaagacatat tgcctcaagt
	2581 tggcaaaaat tgaggtgga attgaatgt ggtcaccttg aatggtttg atttaagaaa
	2641 aaatagataa ctgtattgt aaatatctt aaaatatctt tattcatcc ctgagaaatt
25	2701 tgtgtggtat gtctgattg ctctcccag atctgcctt gtctttact cacacaact
	2761 tgtgtcttt ttgtaaagaa acaaaacaag agccatgcac accagttgt gctctcaaa
	2821 tgtactcagc tgtgtggcca tctgtgggt tctggtgcc ttaccagggg ctacattct
	2881 ggagaacact gccttcctt ttcccccacc acctattgt aattgtctt catgtccagc
	2941 ttctctctc ttgctggat ttggtctgac ttgggctgc acggtcgggt gcaggctgtc
30	3001 agaagcgtg tgaagatagc tcgggtagt taagtctacc tcaggcatc caacaaggcc
	3061 ctcaaatga ggctttgct ttctgtgtc tcttagtgag tgatatact attctaactg
	3121 gctattcata catttcatc agtgtgggc aataaatggg acaatttaa ggagcctcaa
	3181 ttctaagac tggttattc caccagggc ttgatatgg ttgacctgcc ttgccaacag
	3241 gtgcaagtat catatatgc agtgcctgg ttgaaatgt gtgtgtgtgt gtgtgtgtgt
35	3301 ccgtgtgtgt gtgtgtgtgt gtgtgtgtgt aaggagggt ggaagggtga tgggtggaga
	3361 caggaaattc cagatgtga gatttcagt tagaaattat atgtgtgtgt gtgtgtgtct
	3421 gtctgtctgt ctggactta ttgcaggta cttccagga ccagggtacc ccagttcaca
	3481 ctgggtttag agttgccaag ctcaagtata agcttggtt ggtagacaga tggcctcac

3541 ctcaactcct ggccctgggg cttgtctca aggcacctca ttttagttg tagaataatt  
 3601 gaagggaccc cagctttct tagcttctc ttgacagcta taaggaaggg tgaagcatct  
 3661 ttttcagaga tcttagaatt gtgtctcac ttctgtcaag taataaaca tatatatcca  
 3721 ttgatgttt attctattcc cctattaacc ttggattta atcaaggaca tttatgatg  
 5 3781 tgcaagggtg taatcattaa ttctgtgga aggtcacaag ataggagaaa acaattcttt  
 3841 ctatagtaaa acaccatgat acaataaat ttagttttag aaaatgggaa cctgaagttt  
 3901 tgattcacat agattttat agttttacag gctccattcc aatgtatgaa aaatatgtat  
 3961 ctgattctgt gaatttgcatt tgcaaagggg gaaagatttc actcttgaag cctctctcct  
 4021 tcagctcctc cctcagtcg agactgcata gtgccgggt aagggtgggg tgctctttg  
 10 4081 cctcaggagt gctgttcag cagcaggctc tgcaagggtga cctttgctt gctcagaaga  
 4141 cactgatgat caagatgctg gctgtggctc cgagacctga tgccagtga gaggaagatg  
 4201 gggtagctag gcaactcaa aacagtcaa tggctgcca gcatcgagcg agcggagggt  
 4261 gcacaagctg atgctgtgtg aggaaggag ctaaagatgc ctcagaaag cttttggg  
 4321 gtgattcttc tgccaacccc taggatattg tgagctacag agttattaaa ccagactgag  
 15 4381 gaaacaaaag cccaataaag ctattgaaag tgcccaagct cagagagcag atagcagggg  
 4441 aaggattga attcagggat ctgaaaccaa atcctgtgtt ctctctccta gcctaaact  
 4501 tctctcctt aaacactgta agaggaagat ttctctctt tactgggata acgccaatt  
 4561 ctatatagac cagggtggaa attacaagtg cttatcatt tacaatctac ttttagttaa  
 4621 tgatgcttaa agctagccca ggagagacgt taccctcatg gataacagca tagggccaga  
 20 4681 gccacgagct atgtactctg tatcttcag gctgttgcct ccacaggcag gtagagtcag  
 4741 aagccatgac agtctgagc atgcagaggc cccacatac ccaggttat ttctggaacc  
 4801 tgggtgttt tctcacatta gtacttctc ctgtcctag aaaagggcca aatgaagac  
 4861 caaaatattg gggtagctg gctgtcatct tcatcttat gaccggttt gtggtgttct  
 4921 ttgtctaaa cag (SEQ ID NO:7; nucleotides 1-1627 of GENBANK™ Accession  
 25 No. AJ002141)

Expression of the nucleic acid is preferably inducible. Osteoblast or odontoblast  
 transcriptional regulatory DNA is used to control expression of IL-4 or another anti-  
 inflammatory polypeptide in a transcription unit. A truncated fragment of such promoters,  
 30 e.g., containing part of SEQ ID NO:3, 4, 5, 6, or 7, which functions to preferentially direct  
 transcription in odontoprogenitor cells or OPCs (compared to other cell types) may be used.  
 The regulatory sequence, e.g., a cis-acting cell-specific transcriptional regulatory element, is  
 positioned 5' to a heterologous nucleic acid sequence, in a transcription unit. All or part of  
 one of the nucleotide sequences specified in a reference sequence, its complementary strand  
 35 or a variant thereof may be used in to direct transcription of a heterologous nucleic acid  
 sequence. A nucleic acid fragment is a portion of at least 20 continuous nucleotides identical

to a portion of length equivalent to one of the reference nucleotide sequences or to its complement.

Expression of a heterologous polypeptide-encoding sequence is regulated by contacting the cells of the invention with an antibiotic compound such as tetracycline or a tetracycline analogue (e.g., minocycline or doxycycline). For example, tetracycline is systemically administered at least 2 days before periodontal surgery, e.g., and the time at which cells of the invention are implanted, and/or for at least 2 days after surgery and/or implantation. Expression of the heterologous polypeptide by the cells is turned on while the antibiotic is present in the tissue, i.e., while it is being administered to the cell implant recipient. Expression of the recombinant anti-inflammatory polypeptide decreases and ceases after administration of the antibiotic is stopped. Typically, an antibiotic administered 8-12 days prior to surgery and 8-12 days post-surgery. Similarly, antibiotics are administered before and after orthopedic surgery, e.g., surgery for cartilage removal from articulating joints or for removal of metastatic bone tumors (at which time the cells are implanted at or adjacent site to diseased tissue). The cells may be implanted before, during, or after implantation of a dental orthopedic prosthesis. To treat advanced periodontal disease, the cells are administered locally to the periodontal ligament in the mandibular section of the jaw. A clinical benefit is conferred by using the cells to inhibit osteolysis in a mammal, e.g. a human patient, that is suffering from or at risk of developing periodontitis or other bone disorders which may lead to bone loss, e.g., alveolar bone loss.

The methods described herein are also applicable to veterinary use, e.g., to treat dogs, cats, horses.

The invention includes OPCs which are genetically modified to contain a nucleic acid encoding an anti-inflammatory polypeptide. OPCs are derived from bone marrow stromal cells and have been differentiated *ex vivo* in the presence of ECM. As is described above, the OPCs preferably contain a nucleic acid encoding a cytokine such as IL-4, or an agonist thereof, operably linked to a promoter which directs transcription of a nucleic acid to which it is linked preferentially in cells which have differentiated into osteoblasts.

For treatment of bone disorders, the cells are implanted into the bone marrow of a recipient mammal or into an articulating joint of the mammal. For example, the cells are administered intratibially or intrafemorally. The cells are implanted locally, e.g., at the site

of bone loss or adjacent to such as site, e.g. in the bone marrow, and expression of the recombinant polypeptide by the cells is regulated by systemically administering an antibiotic such as minocycline or doxycycline. Methods of transplanting cells into the bone marrow of a mammal are well known in the art, e.g., as described in U.S. Patent No. 4,188,486. The dose of cells to be administered ranges from  $1 \times 10^5$  cells to  $1 \times 10^{10}$  cells in volume suitable for the location of transplantation (e.g., a smaller volume is used for implantation into mandibular tissue or into the periodontal ligament compared to implantation into the bone marrow of the femur). Clinical protocols for such implantation procedures are known in the art. For example, a dose of  $1 \times 10^8$  cells per kg of body weight is administered to femoral bone marrow. Repeated implants may be required in the case of long term diseases such as rheumatoid arthritis.

Inhibitors of cyclooxygenase II (COX-2) or tumor necrosis factor-alpha (TNF $\alpha$ ) are optionally administered. COX inhibitors include aspirin, ibuprofen and indomethacin, as well as bisaryl COX-2 inhibitory compounds (e.g., as described in U.S. Patent No. 5,994,379) and (methylsulfonyl)phenyl-2-(5H)-furanones (e.g., as described in U.S. Patent No. 6,020,343).

The isolated genetically-modified OPCs are used to treat individuals suffering from or at risk of developing a bone loss disorder such as rheumatoid arthritis, osteoporosis, periapical or endochondral bone loss, artificial joint particle-induced osteolysis, bone fracture or deficiency, primary or secondary hyperparathyroidism, metastatic bone disease, osteolytic bone disease, post-plastic surgery, post-prosthetic joint surgery.

By the term "osteoprogenitor" is meant a differentiated bone precursor cell derived from a bone stromal cell. By the term "odontoprogenitor" is meant a differentiated bone precursor cell derived from periodontal ligament. The differentiated state of the bone marrow stromal cells or ligament derived cells is induced by culture in the presence of ECM. Preferably, the cells are cultured in the presence of a BMP such as BMP-2, 4, or 6. Differentiated progenitor cells have enhanced ability to build bone tissue, compared to undifferentiated stromal cells. OPCs or odontoprogenitor cells are distinguished from bone stromal cells (as well as fat, muscle, or cartilage cells or tissue) by the production of alkaline phosphatase, expression of osteocalcin, and expression of bone sialoprotein (in addition to the expression of dentin sialoprotein in the case of odontoprogenitors).

The *ex vivo* cell-based therapeutic methods of the invention has several advantages over standard gene therapy protocols. For example, the cells expressing the recombinant anti-inflammatory polypeptide are isolated, i.e., purified from cells which do not have the desired phenotype. A population of isolated OPCs or odontoprogenitor cells is at least 75%,  
 5 more preferably 85%, more preferably 90%, more preferably 95%, more preferably 98%, more preferably 99% or 100% OPCs or odontoprogenitor cells, respectively.

DNA is introduced into isolated cells *ex vivo*, thus avoiding or minimizing the possibility DNA uptake by non-target cells in the body. Another measure of safety is conferred by using a transcriptional regulatory element and a promoter that directs  
 10 transcription only in the isolated cell type. *In vivo* expression of the recombinant polypeptide is further regulated by the systemic administration of an antibiotic or antibiotic analogue.

OPCs are isolated and expanded from stromal cells from bone marrow aspirates, and autologous bone marrow stromal cells are expanded. The cells are optionally frozen and stored in liquid nitrogen for long periods of time before being differentiated and transduced.  
 15 These "banked" autologous cells allow for multiple inoculations over a long period of time, which is advantageous since RA may persist for many years

Other features and advantages of the invention will be apparent from the following detailed description and from the claims.

#### BRIEF DESCRIPTION OF THE DRAWINGS

20 Fig. 1 is a bar graph showing alkaline phosphatase activity of rabbit bone marrow stromal cells plated on ECM produced by untransduced C3H10t1/2 cells, or C3H10T1/2 cells transduced with and without BMP-6, or on plastic alone. Activity is expressed as  $\mu\text{mol p-nitrophenol produced/min/mg protein} \times 10^{-4}$  at days 1 and 21.

Fig. 2A is a diagram of a periodontal ligament biopsy.

25 Fig. 2B is a diagram of the method for differentiating odontoblast precursor cells on an osteoinductive matrix and transduction of the cells with a regulatable therapeutic gene.

Fig. 2C is a diagram of a device for implanting cells adjacent to teeth in need of therapeutic intervention.

30 Fig. 3A is a photomicrograph of rabbit bone marrow stromal cells cultured for 21 days after plating on tissue culture plastic. Magnification, 100 $\times$ .

Fig. 3B is a photomicrograph of rabbit bone marrow stromal cells cultured for 21 days after plating on ECM-coated dishes from a C3H10T1/2 cell line transduced with rhuBMP-6. Magnification, 100×.

Fig. 4 is a bar graph showing the effect of rhuIL-4 on rhuIL-1 alpha-induced PGE2 by rabbit osteoprogenitor cells.

#### DETAILED DESCRIPTION

Cell-based delivery of rhIL-4 at the site of an osteolytic lesion allows rhIL-4 to be concentrated near an inflammatory site where inflammatory effector cells, e.g., macrophages, and osteolytic effector cells, e.g., osteoclast precursor cells, are located. Adverse effects of rhIL-4 on thymocyte and T cell function are greatly decreased since the cytokine acts locally as opposed to functioning throughout the body when delivered systemically.

The cells described herein are committed to the osteoblastic lineage. Differentiation was induced by stimulating the bone marrow stromal cells to differentiate by exposing the cells to an extracellular matrix such as Matrigel (Becton Dickenson) or other commercially available matrix preparation in the presence of bone morphogenetic proteins. This step is typically carried out before the differentiated cells are transduced with retroviral expression vectors containing genes encoding one or more therapeutic proteins. This is advantageous in that the transduced cells cannot differentiate into cartilage, muscle or fat cells once implanted *in vivo*. In contrast, the pre-osteoblastic pluripotent bone marrow stromal cells still retain the potential to differentiate.

Unlike the retroviral vectors employed in the prior art which use viral promoters, the retroviral expression vectors in the OPCs have been constructed to use osteoblast specific promoters to initiate transcription of the reverse transactivator Tetracycline Activator (rtTA) gene, which in turn regulates production of the rhIL-4 therapeutic protein. This approach provides the advantage of increased safety because the osteoblast promoters direct transcription more efficiently in OPCs (compared to other cell types such as stromal cells) and are far less likely to be inactivated *in vivo* than viral promoters. For example, the viral vectors contain a doxycycline-inducible system which regulates the expression of the interleukin-4 encoding sequences.

The OPCs described herein were modified to increase the expression of the alpha-5 integrin receptor. This modification allows the cells to adhere to bone matrix proteins when implanted *in vivo*, which confers the added advantage that the OPCs may be inserted directly

into osteolytic sites without prior encapsulation, e.g., in porous calcium phosphate ceramic cubes or other types of encapsulated devices.

The cells are used for autologous or allogeneic cell transplants to serve as a cell-based platform to deliver the recombinant human interleukin-4 (rhIL-4) protein in a site-specific, regulated manner. RhIL-4 acts on defined cell types when the OPCs are implanted at the site of osteolytic bone lesions. The OPCs, genetically-engineered to produce rhIL-4 are used to i) promote tumor reduction when used in conjunction with anti-cancer drugs, ii) inhibit formation of osteoclasts which resorb bone, and iii) stimulate new bone growth. The methods result in improved clinical outcomes. The OPCs, engineered to secrete rhIL-4 may are implanted in patients who are undergoing revision of a artificial joint replacement due to the development of implant-induced osteolysis, as well as in patients suffering bone loss due to rheumatoid arthritis and in the oral cavity due to severe periodontal disease.

#### IL-4 and agonists thereof

IL-4, an anti-inflammatory cytokine produced primarily by Th2 cells and macrophages exhibits anti-inflammatory and immunosuppressive properties. Bone-derived cells, e.g., differentiated osteoprogenitor cells or odontoprogenitor cells, which are genetically-altered to produce recombinant human interleukin-4 (rhuIL-4) are administered to diseased bone tissue. The invention provides a gene therapy approach to deliver rhuIL-4 locally at inflamed joints by targeting bone marrow stromal cells that have undergone partial differentiation with a viral plasmid expression system containing a cell-specific promoter. In order to prevent potentially harmful effects of rhuIL-4, due to high local or systemic concentrations, the production of rhuIL-4 is regulated by the oral administration of antibiotic analogues. Articular cartilage degradation and bone resorption, associated with rheumatoid arthritis, is reduced significantly by the local, regulated release of IL-4 near the site of tissue damage. The mechanism for reduced cartilage degradation and bone loss is based on the ability of IL-4 to inhibit TNF-alpha, IL-1, and PGE2 production, as well as the ability of IL-4 to decrease angiogenesis. Regulated, local release of IL-4 decreases cartilage and bone destruction *in vivo*. IL-4 also has immunosuppressive properties; in situations in which the implanted cells are allogeneic (rather than autologous), the IL-4 produced by the implanted cells may obviate the need to administer systemic immunosuppressive drugs to combat tissue rejection.



Example 1: Differentiation of rabbit bone marrow stromal cells on a BMP-6 containing osteoinductive extracellular matrix derived from C3H10T1/2 cells

Over 185,000 spinal arthrodeses are performed in the US each year, with non-union rates as high as 35% reported in the most commonly performed procedure, posterolateral lumbar intertransverse process fusions. Autologous iliac crest bone is the gold standard graft material, but it is limited in quantity, and the morbidity of harvest is not insignificant. The data described herein indicate that BMPs induce differentiation of multipotential stromal cells from rabbit bone marrow an osteoblastic lineage. The stromal cells are exposed to ECM secreted by a transfected murine cell line constitutively expressing or overexpressing mRNA for BMP-6.

Cells from the C3H10T1/2 murine fibroblast line were transduced with either an LXS vector containing the rhBMP-6 gene or the same vector without the gene. The cells were cultured under standard conditions in DMEM (Gibco; Gaithersburg, MD) supplemented with 100 units/ml PCN, 100 µg/ml streptomycin, and 10% FBS (Hyclone; Logan, UT). Four days after reaching confluence, the cells were lysed sequentially with water and 0.1% Triton X-100. The plates were gently washed with phosphate buffered saline (PBS), leaving the extracellular matrix and the BMP-6 protein adherent to the plastic dish.

Bone marrow was aspirated from the femurs of two New Zealand white rabbits and suspended in DMEM with 100u heparin/cc. The cell suspension was diluted with PBS, 2%BSA, 0.6% sodium citrate, and 1% penicillin/streptomycin. The suspension was then layered on a Ficoll-Paque (Amersham Pharmacia Biotech; Piscataway, NJ) gradient, and centrifuged at 600 × g for 20 minutes. The cells at the interface were isolated, washed, and re-centrifuged at 500 × g twice. They were then cultured to confluence in a T-75 flask under standard conditions in a-MEM with L-glutamine 2mM, without nucleosides, supplemented by 12.5% FBS, 0.2mM I-inositol, 20nM folic acid, 0.1mM beta-mercaptoethanol, and 1% penicillin/streptomycin. The cells were re-plated in triplicate on the ECM produced by untransduced cells, LXS transduced cells, or LXS-BMP6 transduced cells, or on plastic, and cultured for 21 days.

Alkaline Phosphatase (ALP) activity was determined on day 1 and 21 after plating of stromal cells on ECM. The plates were scraped and rinsed with 0.5M CAPS, pH 10.5, and

sonicated. 0.5ml of 0.5% p-nitrophenyl phosphate was added to each sonicate, and incubated at 37° C for 30 minutes. 0.2M NaOH was added to stop the reaction, and the amount of p-nitrophenol produced was determined by spectrophotometry at 405nm. ALP activity was expressed as  $\mu\text{mol p-nitrophenol/min/mg protein}$ . Protein content was determined by  
5 Bradford protein assay (Bio-Rad, Hercules, CA). As an additional control, the ECM produced by each C3H10T1/3 cell line was also assayed for ALP activity at day one, and after exposure to media for 21 days.

The ALP activity in the ECM alone was negligible at day 1 and 21. The activity from each of the ECM-exposed marrow cells was likewise negligible at day 1, as was the activity  
10 from the cells on plastic. However, at day 21, there were striking differences (Fig. 1). The ALP level produced by marrow cells plated on plastic was unchanged. That of marrow cells plated on the ECM from untransduced or LXS cells each increased 400%, while that of the marrow cells plated on the ECM from BMP-6 transfected cells increased 700%.

ECM-bound BMP 2 and 4 produced by neonatal mouse calvarial cells stimulated  
15 ALP activity in mouse bone marrow cells. Exposure of stromal cells to ECM in the absence of BMP-6 increased ALP production, presumably due to the presence of type collagen in the matrix. The further increase of ALP production by BMP exposed cells is due to increased osteoblastic differentiation of the stromal cells.

These results indicate that exposure to ECM-bound BMPs induces stromal cells to  
20 differentiate along an osteoblastic lineage. Cells of the osteoblastic lineage, e.g., OPC's or odontoprogenitor cells, are identified and purified by virtue of their expression of marker genes such as alkaline phosphatase, osteocalcin, and bone sialoprotein (in addition to dentin sialoprotein in the case of odontoprogenitors). Probes to detect the marker genes are known in the art (e.g., as described by Guo et al., 2000, *Calcified Tissue International* 66:212-216).  
25 Marker gene expression is detected by measuring transcription of the genes (e.g., using labeled nucleic acid probes in *in situ* assays) or by immunohistochemistry to detect antibody binding to the gene products. The assays described above are used to distinguish stromal cells from OPCs and odontoprogenitor cells.

Example 2: Inhibition of Alveolar Bone Loss by Cell-Delivered IL-4

The osteoclast is responsible for mediating excessive bone resorption during progressive periodontitis. IL-4 inhibits osteoclast differentiation and function. Autologous cells are engineered to express IL-4 and permanently implanted at sites of inflammation, e.g., in the mandible, in soft tissue adjacent to affected teeth, or in the periodontal ligament, using methods known in the art.

Periodontal disease is induced in C3H mice by repeated injections of LPS derived from the clinically-relevant microorganism *Porphyromonas gingivalis*, an art recognized model of periodontal disease. Mice with periodontal disease are treated using C3H10T1/2 cells genetically engineered to produce IL-4 in a regulatable manner. Production of interleukin-4 is regulated by providing antibiotic orally, e.g., in the drinking water. Cells are implanted locally, at sites of bone resorption, thereby bypassing the need for either systemic administration or repetitive local injections of a bioactive molecule. Optionally, antibiotics are placed in the periodontal pocket following implantation of cells for periodontal disease. This cell-based approach for local delivery of interleukin-4 utilizes tissue engineering to inhibit resorption of alveolar bone.

The murine molony retroviral vectors used herein are well characterized and are non-immunogenic in humans or mice.

Standard *in situ* hybridization (ISH) is used to detect IL-4 production as well as characterize the osteoclast phenotype in cells that have populated mandibler bone or other bone tissue of cell implant recipients.

Example 3: Gene therapy vector

A tetracycline analog-regulated expression system is used to direct production of recombinant anti-inflammatory compositions. A dual "tet-on" retroviral system is used for the following reasons; i) the vectors are commercially available and the packaging cells produce high retroviral titers, ii) the murine molony retroviral vectors have been well characterized, are non-immunogenic, and have been used in safely in humans, and iii) the use of two retroviral vectors in the "tet-on" mode prevents "leaking", i.e., recombinant polypeptide expression is extremely low or absent without antibiotic present.

"Tet-off" and tet-on" systems use the antibiotic tetracycline or various analogues to regulate expression. Toxicity of the VP16 viral transactivator fusion protein (tTA) was not

observed and no antibodies were made to the “reverse” (rtTA, tet-on) or rTA, “tet-off” transactivator. The tetracycline analogue doxycycline is the preferred antibiotic inducer for the “tet-on” system and is administered by orally or by intraperitoneal administration using known methods, e.g., as described for mifepristone and rapamycin, as well as by

5 implantation of subcutaneous pellets. Doxycycline and/or minocycline is given orally. Minocycline inhibits the action of matrix metalloproteinases (MMPs) which are involved in breakdown of bone and cartilage. Minocycline-activated site-specific IL-4 production at the inflamed joint acts in a synergistic manner to inhibit inflammation and angiogenesis. These two drugs may work together and lead to increased benefit for patients suffering from

10 RA. The ability to shut down local IL-4 production by removal of minocycline in the “tet on” gene system is advantageous to prevent deleterious effects of sustained IL-4 production at a site of inflammation such as in a rheumatic joint.

Some viral promoters/ enhancers used in adenoviral and retroviral plasmid vectors are inactivated by interferon-gamma and tumor necrosis factor-alpha *in vivo*. These include the

15 rous sarcoma virus (RSV), simian virus 40 (SV-40), and cytomegalo-virus (CMV) promoters. Since levels of IFN and TNF are elevated in RA and OA patients, the use of these standard viral vectors could limit recombinant polypeptide expression, especially if sustained production is required. Given these limitations of viral promoters for long term *in vivo* use, the invention utilizes a constitutive cellular promoter in place of the CMV promoter

20 to control expression of the rtTA transactivator in one of the two retroviral plasmid vectors. A human osteocalcin promoter sequence (e.g., SEQ ID NO:3) is employed to modify a “tet on” retroviral vector for transduction of OPCs *ex vivo*.

Late-stage rabbit osteoprogenitor cells obtained from bone marrow stromal cells were isolated and characterized using marker gene detection. The rabbit OPCs have undergone

25 partial differentiation on a osteoinductive matrix derived from C3H10T1/2 cells that have been transduced with a retroviral vector expressing recombinant human bone morphogenetic protein-6 (rhuBMP-6).

Bone marrow stromal cells are obtained from an individual such as a human patient 8-12 weeks prior to therapy. The cells are expanded, differentiated and transduced with

30 recombinant DNA encoding anti-inflammatory polypeptides *ex vivo*. The OPCs are implanted in the marrow bones adjacent to the diseased or injured site, e.g., in the mandible

or periodontal ligament for periodontal disease or in the marrow of the distal femur and proximal tibia, i.e., in juxtaposition to an inflamed knee joint. This approach allows the rhuIL-4-transduced OPCs to be in close proximity to the bone-resorbing osteoclasts and the pannus/bone interface of the joint. Correct positioning of the implanted OPCs in the marrow is important for the following reasons; i) inhibition of bone resorption by IL-4 is optimized by close proximity of the OPCs to the synovial fibroblasts of the invading pannus and the surrounding osteoblasts, ii) the locally produced IL-4 inhibits osteoclast formation from the differentiated synovial marrow-derived macrophages in the presence of rheumatoid synovial fibroblasts, and iii) the OPC-produced IL-4 locally inhibits neovascularization of the inflamed joint.

Other advantages of using bone marrow stromal cells, which have been partially-differentiated toward the osteoblastic lineage, include a high level of expression of the rhuIL-4 for a given cell population due to the higher number of osteoblastic cells locally present. Moreover, a pure population of differentiated stromal cells is not required due to the high specificity of the osteocalcin promoter for osteoblasts. Adipocyte, muscle, and chondrocytic precursor cells will not express the rhuIL-4 even if these cell types take up the recombinant anti-inflammatory polypeptide-encoding DNA. These features provide a safer and more controlled environment for cytokine release when the transduced osteoblast cells are inoculated into the bone marrow.

#### Example 4: Construction of the rhIL-4 "tet-on" dual retroviral expression vectors

The pRevTet-On and pRev-TRE retroviral expression vectors are publically available from Clontech Laboratories, Inc. (Palo Alto, CA). Both vectors were derived from pLNCX, a retroviral vector capable of producing high-titer virus in the Retropak™ (Clontech, Inc.) PT67 packaging cell line. The core murine Moloney leukemia viral vector for each plasmid consists of a 5' long terminal repeat (LTR) containing a promoter (L) which drives the extended retroviral packaging signal. The remaining DNA consists of pBR322-based plasmid sequences which allow for replication in bacteria and an ampicillin resistance gene for bacterial selection. All plasmids and retroviral expression vectors were purified by cesium chloride/ethidium bromide ultracentrifugation gradients, checked for purity on agarose gels, and analyzed for orientation by restriction enzyme mapping and DNA sequence analysis.

pRev-Tet-On-huOC plasmid construct

The pRevTet-On vector also contained a neomycin phosphotransferase gene, then an internal minimal immediate early cytomegaloviral (CMV) promoter which drives the reverse tetracycline regulatory element (rtTA). A 1.339 kb BamHI/EcoRI cDNA was excised from the pGoscas vector which contains a cDNA of the human osteocalcin promoter. The regulatory DNA was ligated into pRevTet-On digested with BamHI/ClaI, which removes the CMV promoter and the rtTA portion of the vector. A 1.05 kb cDNA to rtTA was subsequently ligated into the ClaI site of pRev-Tet-On followed by blunt end ligation. This resulted in substitution of the viral CMV promoter/enhancer for the osteoblast specific human osteocalcin promoter in the pRev-Tet-On retroviral vector.

pRev-TRE-rhuIL-4 plasmid construct

The pRev-TRE vector (Clontech), contains a 5' long terminal repeat (LTR) containing a promoter which drives the extended retroviral packaging signal. The transactivator response element (TRE) contains seven direct repeats of the tetO operator sequence upstream of a minimal CMV promoter, which can be bound by the tTA and rtTA transactivators. The rhuIL-4-pCD plasmid was obtained from American Type Culture Collection (ATCC (#57593)). A 0.86 kb BamHI hIL-4 insert was isolated and used for subsequent subcloning into the BamHI site of the multiple cloning site in the pRev-TRE vector. Restriction enzyme mapping was performed to check for the correct orientation of the rhuIL-4 cDNA insert. The pRev-TRE-rhuIL-4 vector was subsequently transfected into the PT67 packaging cell line (Clontech), selected with Hygromycin B, and high titer clones were assayed using serial dilutions of viral supernatants before infection and Hygro B selection of NIH3T3 cells. PT67 packaging cells containing the pRev-TRE-rhuIL-4 vector were used to sequentially infect the rabbit osteoprogenitor cells along with the pRevTet-On-huOC vector. The transduced cells are subjected to selection with G418 and Hygromycin B and clones producing rhuIL-4 in response to tetracycline ana-log treatment are isolated. The optimal doxycycline and minocycline concentrations for induction of rhuIL-4 production in the transduced cells is performed using standard methods.

Example 5: Isolation and Osteogenic Differentiation of Rabbit Bone Marrow Stromal Cells

Bone marrow aspirate (approximately 1-2 c.c.) was obtained from the greater trochanteric region of the femur of anesthetized New Zealand white rabbits and suspended in DMEM

with 100u heparin/cc. The cell suspension was diluted with PBS, 2%BSA, 0.6% sodium citrate, and 1% penicillin/ strep-tomycin. The suspension was then layered on a Ficoll-Paque (Amersham Pharmacia Biotech; Piscataway, NJ) gradient, and centrifuged at  $600 \times g$  for 20 minutes. The cells at the interface were isolated, washed, and recentrifuged at  $500 \times g$  twice.

5 They were then cultured for 2-3 weeks until confluent in T-75 flasks at 37 degrees C in 5% CO<sub>2</sub> in alpha-MEM with L-glutamine 2mM, without nucleosides or glucocorticoids, supplemented by 12.5% horse serum (Sigma; St. Louis, MO), 12.5% FBS, 0.2mM I-inositol, 20nM folic acid, 0.1mM B-mercaptoethanol, and 1% penicillin/streptomycin. The cells were then harvested in DPBS/EDTA /pancreatin and stored in liquid nitrogen in freezing media.

10 Aliquots of BMSCs were subsequently thawed and replated in triplicate on plastic dishes or dishes coated with the ECM produced by untransduced cells, LXS transduced cells, or rhuBMP6 transduced cells, and cultured for 1 and 21 days. Examination of the thawed cells by phase contrast microscopy revealed that BMSCs cultured on tissue culture plastic for 21 days retained their fibroblastic, spindle-shaped morphology, as shown in Fig. 3A. In contrast, BMSCs plated on the BMP-6-containing ECM for 21 days became more  
15 cobblestone-shaped in appearance (Fig. 3B) and resembled osteoblast cells.

Alkaline phosphatase (ALP) activity was determined on day 1 and 21 after plating of stromal cells on ECM. Representative results are shown in Fig. 1. These data indicate that stromal cells may be isolated, expanded, and stored frozen before undergoing further  
20 differentiation *ex vivo*. The data also indicate that BMP-responsiveness with respect to alkaline phosphatase induction demonstrates that the BMSCs have undergone partial differentiation to become late stage osteoprogenitor cells

#### Example 6: Effects of rhuIL-4 on Rabbit late-stage Osteoprogenitor Cells

The effect of rhuIL-4 on PGE2 production by late-stage rabbit OPCs was analyzed.  
25 OPCs were harvested from the BMP-6/ECM coated dishes by trypsinization, counted, then replated in 6-well dishes in alpha-MEM plus 1% FBS and  $10^{-5}$  M arachidonic acid for 24 hr. The cells were then incubated for an additional 24 hr with rhuIL-1 (2 ng/ml) in the absence or presence of IL-4 (25, 50, and 100 ng/ml). PGE2 levels were measured in the conditioned cell cul-ture media by enzyme immunoassay (BioTrak RPN 222, Amersham Pharmacia  
30 Biotech, Inc., Piscataway, NJ). As shown in Fig. 4, there was dose-related effect of IL-4 on the inhibition of IL-1-stimulated PGE2 release by the rabbit OPCs.

These results indicate that rhuIL-4 blocks the effect of IL-1-alpha on the induction of PGE2 production in osteoprogenitor cells, an important intermediate step in osteoclast-mediated bone resorption and that IL-4 is beneficial for reducing inflammation associated with RA (since PGE2 is a potent mediator of the pain and edema associated with rheumatoid synovitis).

Other embodiments are within the following claims.

What is claimed is:



1 1. An isolated odontoprogenitor cell comprising a nucleic acid encoding an anti-  
2 inflammatory polypeptide

1 2. The cell of claim 1, wherein said cell is derived from a periodontal ligament..

1 3. The cell of claim 1, wherein said polypeptide is a cytokine.

1 4. The cell of claim 3, wherein said cytokine is interleukin-4 (IL-4).

1 5. The cell of claim 1, wherein said nucleic acid is operably linked to an  
2 osteoblast-specific promoter.

1 6. The cell of claim 5, wherein said osteoblast-specific promoter is an  
2 osteocalcin promoter.

1 7. The cell of claim 5, wherein said osteoblast-specific promoter is a bone  
2 sialoprotein promoter.

1 8. The cell of claim 1, wherein expression of said nucleic acid is inducible.

1 9. The cell of claim 1, wherein expression of said nucleic acid is regulated by an  
2 antibiotic compound.

1 10. The cell of claim 9, wherein said antibiotic compound is tetracycline or a  
2 tetracycline analogue.

1 11. The cell of claim 10, wherein said tetracycline analogue is minocycline or  
2 doxycycline.

1 12. A method of inhibiting osteolysis in a mammal, comprising introducing into  
2 said mammal an isolated odontoprogenitor cell comprising a nucleic acid encoding an  
3 anti-inflammatory polypeptide.

1 13. The method of claim 12, wherein said mammal is suffering from or at risk of  
2 developing periodontitis.

1 14. The method of claim 12, wherein said mammal is suffering from or at risk of  
2 developing alveolar bone loss due to periodontal disease.

1 15. The method of claim 12, wherein said cell is administered to the periodontal  
2 ligament in the mandibular section of the jaw.

1 16. An isolated osteoprogenitor cell comprising a nucleic acid encoding an anti-  
2 inflammatory polypeptide.

1 17. The cell of claim 16, wherein said polypeptide is a cytokine.

1 18. The cell of claim 17, wherein said cytokine is interleukin-4 (IL-4).

1 19. The cell of claim 16, wherein said nucleic acid is operably linked to an  
2 osteoblast-specific promoter.

1 20. The cell of claim 19, wherein said osteoblast-specific promoter is an  
2 osteocalcin promoter.

1 21. The cell of claim 19, wherein said osteoblast-specific promoter is an bone  
2 sialoprotein promoter.

1 22. The cell of claim 16, wherein expression of said nucleic acid is inducible.

1 23. The cell of claim 16, wherein expression of said nucleic acid is regulated by  
2 an antibiotic compound.

1 24. The cell of claim 23, wherein said antibiotic compound is tetracycline or a  
2 tetracycline analogue.

1 25. The cell of claim 24, wherein said tetracycline analogue is minocycline or  
2 doxycycline.

1 26. A method of inhibiting osteolysis in a mammal, comprising introducing into  
2 said mammal an isolated osteoprogenitor cell comprising a nucleic acid encoding an  
3 anti-inflammatory polypeptide.

1 27. The method of claim 26, wherein said cell is implanted into an articulating  
2 joint of said mammal.

1 28. The method of claim 26, wherein said cell is administered intratibially.

1 29. The method of claim 26, wherein said cell is administered intrafemorally.

1 30. The method of claim 26, wherein expression of said polypeptide is regulated  
2 by an antibiotic compound.

1 31. The method of claim 26, wherein said antibiotic compound is tetracycline or a  
2 tetracycline analogue.

1 32. The method of claim 31, further comprising administering minocycline to said  
2 mammal.

1 33. The method of claim 30, wherein said antibiotic compound is administered  
2 systemically.

1 34. The method of claim 26, further comprising administering an inhibitor of  
2 cyclooxygenase II (COX-2).

1 35. The method of claim 26, further comprising administering an inhibitor of  
2 tumor necrosis factor-alpha (TNF $\alpha$ ).

1        36.     The method of claim 26, wherein said mammal is suffering from or at risk of  
2        developing rheumatoid arthritis.

1        37.     The method of claim 26, wherein said mammal is suffering from or at risk of  
2        developing periapical or endochondral bone loss, artificial joint particle-induced  
3        osteolysis, or osteolytic bone metastases.

1        38.     A method of inducing differentiation of a bone marrow stromal cell,  
2        comprising contacting said cell with bone morphogenic protein-6.

**ABSTRACT**

The invention provides compositions and methods to deliver an anti-inflammatory composition, e.g., recombinant human interleukin-4 (rhIL-4), to build (or rebuild) bone tissue. The composition is produced from living osteoprogenitor cells (OPCs) or odontoprogenitor cells.

20006509.doc

20006509.doc




FIG. 1

FIG. 2A




FIG. 2B




FIG. 2C






FIG. 3A




FIG. 3B





FIG. 4



## COMBINED DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled TREATMENT FOR BONE DISORDERS, the specification of which:

- ☒ is attached hereto.  
☐ was filed on \_ as Application Serial No. \_ and was amended on \_\_\_\_\_.  
☐ was described and claimed in PCT International Application No. \_\_\_\_\_ filed on \_\_\_\_\_ and as amended under PCT Article 19 on \_\_\_\_\_.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose all information I know to be material to patentability in accordance with Title 37, Code of Federal Regulations, §1.56.

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose all information I know to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56(a) which became available between the filing date of the prior application and the national or PCT international filing date of this application:

U.S. Serial No.	Filing Date	Status

I hereby appoint the following attorneys and/or agents to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

William E. Booth, Reg. No. 28,933	Ingrid A. Beattie, Reg. No. 42,306
J. Peter Fasse, Reg. No. 32,983	Janis K. Fraser, Reg. No. 34,819
Y. Rocky Tsao, Reg. No. 34,053	Eldora L. Ellison, Reg. No. 39,967
Ruffin B. Cordell, Reg. No. 33,487	

Address all telephone calls to INGRID A. BEATTIE, PH.D., J.D. at telephone number (617) 542-5070.

Address all correspondence to INGRID A. BEATTIE, PH.D., J.D. at:

FISH & RICHARDSON P.C.  
 225 Franklin Street  
 Boston, MA 02110-2804

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patents issued thereon.

04930-028001

**Combined Declaration and Power of Attorney**  
Page 2 of 2 Pages

Full Name of Inventor: Hugh S. Keeping

Inventor's Signature: \_\_\_\_\_ Date: \_\_\_\_\_  
Residence Address: 10 King Philip Avenue, Bristol, RI 02809  
Citizenship: Canada  
Post Office Address: Same as above

Full Name of Inventor: Jonathan S. Reichner

Inventor's Signature: \_\_\_\_\_ Date: \_\_\_\_\_  
Residence Address: 4 Patricia Drive, North Providence, RI 02904  
Citizenship: USA  
Post Office Address: Same as above

20028931.doc

20028931.doc